

Sinbar[®]

herbicide

Season long weed control is easier with Sinbar[®]

Sinbar provides effective residual weed control in apples, peaches, lucerne seed and peppermint.

Features

- Contains 800 g/kg terbacil
- Group 5 mode of action herbicide
- WP formulation (wettable powder)
- Available in a convenient 2 kg pack size that treats between 0.4 and 2 hectares

How Sinbar Works

Sinbar is applied to bare soil under trees, seed lucerne or peppermint crops (as directed on the product label). Rainfall or irrigation is required to move Sinbar into the weed root zone where it is taken up and moved to the leaves. Sinbar inhibits the process of photosynthesis, causing the weeds to yellow and eventually die.

KEY PRODUCT BENEFITS

- Provides season long control of a broad spectrum of grass and broadleaf weeds
- Reduces application costs compared with applying multiple knockdown herbicides
- Controls weeds which are naturally tolerant to, or have developed a tolerance to knockdown herbicides including glyphosate
- Reduces alternative hosts for pests and diseases that may affect crop health
- Increases crop yield potential
- Improves water use efficiency by removing weeds as they germinate, compared to knockdown herbicides which are often applied several weeks after germination
- Has a high level of crop safety when used as directed on the product label

Key Weeds Controlled

Sinbar controls broadleaf and grass weeds including:

- | | | |
|-------------------|--------------------|-------------------|
| • Amaranthus | • Fumitory | • Soursob |
| • Annual ryegrass | • Innocent weed | • Sowthistle |
| • Barley grass | • Johnson grass | • Spiny emex |
| • Barnyard grass | • Kikuyu | • Storksbill |
| • Bathurst burr | • Milk thistle | • Wild lettuce |
| • Brome grasses | • Mustard | • Wild oats |
| • Capeweed | • Nutgrass | • Wild radish |
| • Charlock | • Paterson's curse | • Wild turnip |
| • Chickweed | • Pigweed | • Wireweed |
| • Couch grass | • Sheepweed | • Yellow burrweed |
| • Fat hen | • Shepherd's purse | |

Application Tips

Rates

As a general rule, Sinbar should be used at the lower rates on sandier soils and the higher rates on soils with higher clay content or higher organic matter. Sinbar use should be avoided on very sandy or gravelly soil as crop damage may occur.

See the directions for use table for detailed rate advice.



Timing

Best results are obtained when Sinbar is applied to moist, bare weed-free soil. Established weeds should be controlled by a registered knockdown herbicide prior to application of Sinbar. Rainfall or irrigation 3 to 4 days after application will improve results by moving the herbicide into the root zone.

Crop Safety

Sinbar should be applied only under apples or peach trees that are older than three years.

Sinbar should be applied only on seed lucerne crops that are at least 1 year old.

Do not apply Sinbar to exposed tree roots or crop suckers.

Water Volume

Mix with a minimum water volume of 20 L for every kilogram of Sinbar. Use equipment designed to deliver MEDIUM to COARSE droplet sizes*.

For boom application, use 250–450 L of water per hectare at 200–350 kPa or as recommended by the nozzle manufacturer.

Handguns should deliver 1000–2000 L per hectare at 600–800 kPa.

Directions For Use

The following is an extract of the product label and does not constitute the complete directions for use. The product label should be read thoroughly before opening the packaging.

| CROP | WEEDS CONTROLLED | STATES | RATE kg/ha | | | CRITICAL COMMENTS |
|--|---|-----------------------------|-------------------------------|----------------------------|---|--|
| | | | Light to medium soils | Heavy Soils | | |
| APPLES & PEARS At least three (3) years old | ANNUAL WEEDS: Amaranthus, Barley grass, Barnyard grass, Bathurst burr, Brome grasses, Capeweed, Fat hen, Innocent weed, Milk thistle, Paterson's curse, Pigweed, Shepherd's purse, Threecornered jack(s), (Spiny emex, Doublegee), Whorled pigeon grass, Wild lettuce, Wild oats, Wild radish, Wild turnip | Qld, NSW, Vic, Tas Only | 2.2 kg 3.5 kg | 4.5 kg 4.5 kg | | Apply to moist soil just before or during active weed growth. Light Weed Infestation Heavy Weed Infestation |
| PEACHES At least three (3) years old | As above | SA Only | 2.2–3.5 kg | | | Use higher rates for heavy weed infestation. |
| APPLES & PEACHES At least eight (8) years old | PERENNIAL WEEDS: Couch grass, Kikuyu, Johnson grass, Nutgrass | Qld, NSW, Vic, Tas, SA Only | 4.5 kg | | | Apply as a broadcast spray. Spot Treatment: Complete coverage of weeds is essential. DO NOT apply on sandy or gravelly soils. Note: Where limited rainfall occurs during the growing period, satisfactory control of hard to kill, deep rooted, perennial weeds will not occur. If weeds are present and cannot be removed, add a surfactant/wetting agent*. |
| | | | 200 g/100 L water* | | | |
| LUCERNE SEED CROPS At least one (1) year old | Annual ryegrass, Charlock, Chickweed, Deadnettle, Fat hen, Fumitory, Melilotus, Milk thistle, Mustard, Rough poppy, Paterson's curse, Sheepweed, Shepherd's purse, Soursob, Storksbill (Geranium), Threecornered jack(s) (Spiny emex, Doublegee), Wild oats, Wild radish, Wild turnip, Wireweed, Yellow burrweed | SA Only | Light Sandy Soils 1 kg | Medium Soils 1.4 kg | Heavy High Organic Content Soils 1.75 kg | Apply any time after lucerne becomes dormant, but before growth begins in spring. For best results apply just before the spring germination of weeds. DO NOT use a surfactant/wetting agent. |
| PEPPERMINT | Amaranthus, Barley grass, Brome grasses, Barnyard grass, Capeweed, Fat-hen, Shepherd's purse, Sowthistle, Whorled pigeon grass, Wild oats, Wild radish, Wild turnip | Tas Only | 2 kg | | | Apply during the dormant stage, July to August. |
| | | NSW, Vic Only | | | | Apply 2 to 3 weeks after planting in April–May or to existing plants between the months of April and September. |

Adjuvants and Tank Mixes

A non-ionic surfactant may be added if perennial weeds are established and cannot be removed through other practices. Do not add a surfactant when treating lucerne seed crops.

Tank Mix Compatibility

Overseas experience indicates physical compatibility with diuron, metribuzin, hexazinone and paraquat. As conditions and formulations are rarely consistent, small scale testing should be completed before commercial areas are treated.

Stress

Sinbar should not be applied when weeds are under stress by moisture, temperature, insect damage etc.

Re-cropping Intervals

New crops should not be planted until at least 2 years after the last application of Sinbar herbicide.



*According to specifications by American Society of Agricultural Engineers (A.S.A.E.).

Information and data referred to herein were generated using products and rates that were registered at the time, may include extracts from the product label and does not constitute the complete directions for use. Always read and follow product labels.

AgNova Technologies Pty Ltd shall not be liable for any consequential or other loss or damage relating to the supply or subsequent handling or use of this product, unless such liability by law cannot be lawfully excluded or limited. All warranties, conditions or rights implied by statute or other law which may be lawfully excluded are so excluded. Where the liability of AgNova Technologies Pty Ltd for breach of any such statutory warranties and conditions cannot be lawfully excluded but may be limited to it re-supplying the product or an equivalent product or the cost of a product or an equivalent product, then the liability of AgNova Technologies Pty Ltd for any breach of such statutory warranty or condition is so limited.

© Copyright AgNova Technologies 2026. *Sinbar is a registered trademark of Tessenderlo Kerley Inc. SINAGN260302

For more information scan the QR code or contact your local Area Sales Manager

