

# BLADEX<sup>®</sup> 900WG

## Contact and residual control of tough weeds in vegetables

- Clean crops, higher yields, even maturity.
- Good contact activity plus 4 to 10 weeks control of a wide range of broad leaf and grass weeds including tough weeds like Black Nightshade, Nettles and certain Brassica weeds.
- Good crop safety.
- Short persistence with low potential for plant back issues.

### Mode of Action

Bladex<sup>®</sup>900WG primarily has residual action to control emerging weeds through root uptake. It is also taken up by young emerging weed shoots and has contact action through the foliage of small emerged weeds (<3 cm).

Weed control is primarily due to the active ingredient affecting photosynthesis causing energy starvation.

Bladex needs to be applied to moist soil or have rain or irrigation within 7 days of application to give good residual control.

### Dry flowable formulation

- Easy to Use Dry Flowable.
- Concentrated formula – low use rates.
- Convenient disposable packaging.
- Good compatibility with tank mix partners (always put Bladex in the tank first).
- Improved crop safety in onions over the previous liquid formulation.


### Getting the Best out of Bladex

Do NOT apply Bladex to soils with a high sand or gravel content e.g. sand or sandy loam soils. Bladex is not recommended for use on the Mallee soils of Victoria and South Australia. The activity of Bladex may be reduced in soils of a very high organic matter or clay content.

For best results, weeds should be very young and actively growing. Treatments are more effective on smaller weeds and when the seedbed is free of weeds and large clods.

Best results are obtained if irrigation is applied immediately after spraying. Dry conditions limit root uptake of the herbicide and this may reduce control of all weeds, but particularly grass weeds and wireweed.

*Continued >*



# BLADEX<sup>®</sup> 900WG

## Vegetables

The length of residual control is maximised in vegetables by timing pre-planting applications immediately prior to planting.

Effective weed control may last from 4 to 10 weeks. The length of control depends on many soil characteristics including: moisture, microbial activity, pH and temperature.

Where heavy infestations of annual ryegrass and wireweed are expected Bladex should only be used in a tank mixture with Stomp\* Xtra or trifluralin. Users should refer to the Stomp Xtra or trifluralin labels for advice on crop safety, follow-crop restrictions (etc).

As new varieties of vegetable crops are developed frequently it is important to test Bladex on a small area of the new variety before treating an entire field. In some states, where legislation allows, Bladex may be used at lower than label rates in tank mixes with other herbicides on varieties known to be susceptible to crop damage at the full rate. Temporary crop damage (yellowing) may occur, particularly where Bladex is applied to crops grown on sand over clay (duplex) soils.

## Mixing Bladex 900WG

Half fill the spray tank and commence agitation. Add the required amount of Bladex to the tank and continue to agitate while filling the tank and during spraying. It is critical to add tank mix partners after Bladex is fully dispersed.

## Compatibility

Bladex is compatible with both Stomp Xtra and trifluralin. Users should refer to these labels for advice on crop safety, follow-crop restrictions (etc).

Bladex may be mixed with a knockdown herbicide such as SPRAY.SEED\* or Roundup\* in pre-plant applications where weeds are dense and/or are greater than 3 cm in diameter.

## Weeds controlled

Annual Ryegrass (*Lolium rigidum*)  
Blackberry Nightshade (*Solanum* spp.)  
Black Bindweed (*Polygonum convolvulus*)  
Capeweed (*Arctotheca calendula*)  
Chickweed (*Stellaria media*)  
Clovers (*Trifolium* spp.)  
Cotula (*Cotula* spp.)  
Crassula (*Crassula* spp.)  
Crowfoot/Erodium/Storksbill (*Erodium* spp.)  
Deadnettle (*Lamium amplexicaule*)  
Docks (*Rumex* spp.)  
Fat Hen (*Chenopodium album*)  
Mountain Sorrel (*Calandrinia caulescens*)  
Mouse Ear Chickweed (*Cerastium* spp.)  
Persicaria/Redshank (*Polygonum persicaria*)  
Plantain (*Plantago* spp.)  
Prickly Lettuce (*Lactuca serriola*)  
Rough Poppy (*Papaver hybridum*)  
Sorrel (*Rumex acetosella*) Sowthistle (*Sonchus oleraceus*)  
Spiny Emex/Three Cornered Jack (*Emex australis*)  
Staggerweed (*Stachys arvensis*)  
Stinging Nettle (*Urtica urens*)  
Turnip Weed (*Rapistrum rugosum*)  
Wild Mustard (*Sisymbrium* spp.)  
Wild Turnip (*Brassica tournefortii*)

## Suppression only

Barley grass (*Hordeum leporinum*)  
Fumitory (*Fumaria* spp.)  
Great Brome (*Bromus driandrus*)  
Wild Radish (*Raphanus raphanistrum*)  
Wireweed (*Polygonum aviculare*)

## AgNova Technologies Pty Ltd

PO Box 590, Eltham, Victoria, Australia 3095  
Ph 03 9899 8100

AgNova Technologies Pty Ltd shall not be liable for any consequential or other loss or damage relating to the supply or subsequent handling or use of this product, unless such liability by law cannot be lawfully excluded or limited. All warranties, conditions or rights implied by statute or other law which may be lawfully excluded are so excluded. Where the liability of AgNova Technologies Pty Ltd for breach of any such statutory warranties and conditions cannot be lawfully excluded but may be limited to it re-supplying the product or an equivalent product or the cost of a product or an equivalent product, then the liability of AgNova Technologies Pty Ltd for any breach of such statutory warranty or condition is so limited.

©Copyright AgNova Technologies 2011

\*BLADEX is a registered Trademark of AgNova Technologies Pty Ltd.  
\*STOMP Xtra, ROUNDUP and SPRAYSEED are registered trademarks.