

Innovation. Quality. Solutions.

Pyramin® WG

SELECTIVE HERBICIDE

Pyramin contains 650 g/kg chloridazon (MOA Group C)

Features

- Unique active ingredient
- Broad weed spectrum
- Taken up by both weed roots and leaves
- Good crop safety

Background

Pyramin® WG Selective Herbicide has been in the market for many years and was developed for the control of annual broad-leaved weeds and various grasses in red beet, silver beet and fodder beet.

Mode of Action

Pyramin is a member of the pyridazinone group of herbicides. Pyramin is a herbicide which inhibits photosynthesis at photosystem II. For weed resistance management Pyramin is a Group C herbicide.

Pyramin acts mainly through the roots of weeds which die a few days after they break through the ground. Pyramin remains effective for several weeks after application. No chemical residues remain in the soil after the crop has been harvested.

Use Rate

Refer to table.

Application

Apply Pyramin the day of sowing or as soon as possible after planting, but before the crop emerges. If soil moisture is inadequate for proper beet germination, Pyramin will not give good weed control, in which case irrigate within 10 days.

Irrigation after application however, should not exceed 35 mm per week, particularly during cold weather conditions.

Pyramin should be applied in at least 110 to 220 litres of water per hectare.



Restraint

DO NOT apply Pyramin to sandy soils with low organic matter as crop damage may result.

Re-entry Period

DO NOT enter treated areas for 10 days unless wearing cotton overalls buttoned to the neck and wrist (or equivalent clothing) and chemical resistant gloves. Clothing must be laundered after each day's use.

Summary

- Pyramin has a unique active ingredient which results in absorption in both the roots and leaves of the weed.
- Broad spectrum herbicide with good crop safety levels.

Innovation. Quality. Solutions.

Pyramin® WG

SELECTIVE HERBICIDE

Directions for Use Table

CROP	WEEDS CONTROLLED	STATE	RATE PER HECTARE Medium to Heavy Soils	CRITICAL COMMENTS
Fodder beet, red beet, silver beet	Bitter cress, blackberry nightshade, black bindweed, capeweed, chamomile, charlock, chickweed, common sowthistle, cotula, deadnettle, docks, dwarf nettle, fat hen, field poppy, groundsel, marshmallow, mustard, pale persicaria, penny cress, persicaria, pig weed, potato-weed, scarlet pimpernel, shepherd's purse, sowthistle, speedwell, spurry, stagger weed, wild radish, winter grass	NSW, Vic, Tas, SA, WA only	4.6 to 6.5 kg	Apply at sowing, or as soon as possible after planting, but before crop emerges.
		Qld only	5.5 to 8 kg	Apply at sowing, or as soon as possible after planting, but before crop emerges.
Baby leaf spinach, Baby leaf beet		Vic and Tas only	4.5 kg	Apply after sowing of direct seeded crops, but before crop emergence.

The information provided herein may include extracts from the product label and does not constitute the complete directions for use. READ THE PRODUCT LABEL THOROUGHLY BEFORE OPENING OR USING PYRAMIN

AgNova Technologies Pty Ltd
 ABN 70 097 705 158
 PO Box 2069 Box Hill North
 Victoria 3129 Australia
 Ph 03 9899 8100

Data referred to herein were generated using products and rates that were registered at the time. Always read and follow product labels. AgNova Technologies Pty Ltd shall not be liable for any consequential or other loss or damage relating to the supply or subsequent handling or use of this product, unless such liability by law cannot be lawfully excluded or limited. All warranties, conditions or rights implied by statute or other law which may be lawfully excluded are so excluded. Where the liability of AgNova Technologies Pty Ltd for breach of any such statutory warranties and conditions cannot be lawfully excluded but may be limited to it re-supplying the product or an equivalent product or the cost of a product or an equivalent product, then the liability of AgNova Technologies Pty Ltd for any breach of such statutory warranty or condition is so limited.
 © Copyright AgNova Technologies 2019. PYR191001
 ® Pyramin is a registered trademark of BASF