

Surround® WP

Crop Protectant

Protection you can see.
Performance you can trust.

Higher, earlier yields: Improved growth of young trees

Modern citrus trees have descended from rainforest species originally adapted to growing in shade conditions. In Australia, citrus is grown in regions where high summer temperatures are common. Air temperatures exceeding 28°C can cause trees to shut down, thereby stopping photosynthesis and stunting growth.

Canopy temperature can be reduced by up to 8°C using a Surround® program, which in turn reduces overall stress on the tree. Over the hot summer months, this can mean a significant improvement in growth of young trees which in turn come into production earlier than would otherwise occur.

Ultimately a Surround program provides the opportunity for earlier fruit production which can significantly improve the financial viability of new citrus plantings.

International trials have demonstrated higher, earlier yields following the use of a Surround program (see data overleaf).

Australian trials have confirmed under local conditions that a Surround program improves overall tree vigour, with treated trees often being larger and having improved foliar flushes.

Getting the best out of Surround

Commence applications prior to the first heat event and maintain the Surround film over the hot summer months. Repeat applications will be required to cover new growth and after heavy rain as well. Spray the canopy to 'near drip' coverage (before the point of runoff).

Surround can disrupt activity of some insects. Monitor Surround treated trees for Citrus Scale and take appropriate action if Citrus Scale is found.



Treated with Surround – Improved growth of young trees.
(2003/04 Mildura)




Untreated

Continued over >


Surround® WP

Crop Protectant


Research conducted by Lapointe, McKenzie and Hall, published in the Journal of Economic Entomology in 2006, has confirmed observations in Australian trials and commercial use.




Source: Lapointe, McKenzie and Hall 2006



Two year results of biweekly applications on Valencia citrus trees. S. L. Lapointe, U.S. Horticultural Research Lab, Ft. Pierce, FLORIDA, USA



Source: Lapointe, McKenzie and Hall 2006



Source: Lapointe, McKenzie and Hall 2006

For best results, maintain Surround coverage throughout the sunburn season.

Always read the product label prior to opening the packaging.

AgNova Technologies Pty Ltd
 ABN 70 097 705 158
 PO Box 2069 Box Hill North
 Victoria 3129 Australia
 Ph: 03 9899 8100

AgNova Technologies Pty Ltd shall not be liable for any consequential or other loss or damage relating to the supply or subsequent handling or use of this product, unless such liability by law cannot be lawfully excluded or limited. All warranties, conditions or rights implied by statute or other law which may be lawfully excluded are so excluded. Where the liability of AgNova Technologies Pty Ltd for breach of any such statutory warranties and conditions cannot be lawfully excluded but may be limited to it re-supplying the product or an equivalent product or the cost of a product or an equivalent product, then the liability of AgNova Technologies Pty Ltd for any breach of such statutory warranty or condition is so limited.

©Copyright AgNova Technologies 2015

*Surround is a registered trademark of Tessenlerlo Kerley, Inc. SURTRE150909