

Innovation. Quality. Solutions.



ACTIVE INGREDIENT **700 g/kg methabenzthiazuron**

FEATURES

- Early post-emergence weed control
- Group C Herbicide
- Control of annual broadleaf weed seedlings

BACKGROUND

Tribunil has been in the market for many years and the use rates have been modified by grower experience. The product was originally developed as a stand-alone herbicide. More recently the product has been used in tank mixtures at reduced rates to improve crop safety and maintain adequate weed control. This is partly due to the adoption of new hybrid onion varieties that are more sensitive to herbicides.

MODE OF ACTION

TRIBUNIL is a member of the urea group of herbicides and has the inhibitors of photosynthesis at photosystem II mode of action.

The transfer of electrons from Photosystem II to Photosystem I is essential for the production of photosynthetic energy. A disruption in this process results in the inability for the plant to fix CO₂ and produce the nutrients needed for survival.

THE DOSE

Rate of application depends on the number of leaves present on crop at time of application (see label)

TIMING

Applied once first true onion leaf has emerged and developed.

Rates will depend on the time of year onions are sown (see label).

APPLICATION

Spray actively growing weeds but not if droughted, frosted or waterlogged.

When rain is imminent spray after and not before. Rain within 12-24 hours after spraying may reduce weed control.

Avoid heavy dew and wait until frost has melted. Do not spray crops wet with dew or rain

COMPATIBILITY

Do not add any pesticide or wetting agent to TRIBUNIL for use on onions.

WITHHOLDING PERIODS

Do not sow susceptible crops such as crucifers, cucurbits, lettuce or tomatoes within 18 months after applying more than 2 kg TRIBUNIL /ha in one season

SUMMARY

- Early application from the 1 leaf stage (see product label for details)
- Wide spectrum broadleaf weed control
- Trusted and proven performance and crop safety

