

Innovation. Quality. Solutions.

Zampro®

Fungicide

Chemistry for downy mildew spray programs

Zampro contains 300 g/L ametoctradin and 225 g/L dimethomorph (MOA Group 45/40)

The Product

Zampro® is a unique fungicide for the control of downy mildew in grapes, and contains the active ingredients dimethomorph and ametoctradin.

Dimethomorph is a systemic morpholine fungicide, with the mode of action being inhibition of sterol synthesis which is vital for cell membrane structure and function (Group 40).

Ametoctradin is a non-systemic fungicide that inhibits respiration in the cell mitochondria (Group 45).

Application Rate

Zampro is registered at 80 mL/100 L for dilute use in grapevines.

Directions for Use Table

The following is an extract of the product label and does not constitute the complete directions for use. The product label should be read thoroughly before opening the packaging.

CROP	DISEASE	RATE	CRITICAL COMMENTS
Grapevines	Downy mildew (<i>Plasmopara viticola</i>)	80 mL/100 L	Apply at 7 to 14 day intervals when humid or wet conditions favour infection but before disease is evident. Use the shorter interval when conditions favouring infection are creating a high risk. Apply a maximum of 2 consecutive sprays before changing to an alternative mode of action for at least one application.

WITHHOLDING PERIOD

Grapes grown for domestic wine: Do not harvest for 4 weeks after the last application.

Grapes grown for export wine: One application per season: refer to point 6 below.

Multiple applications: Do not apply after E-L 25 (80% capfall).

Summary

- Zampro is a new product for AgNova Technologies and is manufactured by BASF.
- Zampro has systemic and non-systemic properties, and should be applied as a protectant fungicide.
- Rainfast:** 1 hour.
- Compatibility:** Zampro has been tested in tank mixture with either Flute or Kusabi fungicides. Zampro is widely compatible with other crop protection compounds but local testing is always recommended before large scale use.
- Crop Safety:** Excellent crop safety when used as directed.

Safety

Zampro has an excellent toxicological and environmental profile when used as directed.

Timing

Zampro is ideal for use against downy mildew in grapevines when wet and warm weather conditions combine to provide a threat of an infection period. 10:10:24 provides a guide, i.e. when the temperature is above 10 degrees Celcius and rainfall is above 10 mm during a 24 hour period. Downy mildew symptoms (oilspots) may not appear for two weeks or more after infection.

Zampro has systemic and non-systemic properties, and should be applied as a protectant fungicide.

6. WHP: Dimethomorph application to grapes grown for export wine production in Australia

Zampro or other products containing dimethomorph (e.g. Acrobat*, Downright* or Sphinx*) can be applied up to growth stage E-L 31 (berries pea-size 7 mm in diameter) provided that only ONE application is made per crop with ANY of these products. If more than one application of products containing dimethomorph is applied to the crop the current E-L 25 (80% capfall) restriction must be observed in order to avoid a detectable residue in wine.

- Safety:** Zampro shows an excellent ecotoxicological profile. It is practically non-toxic to birds, mammals, honeybees, earthworms and other soil macro-organisms, or to non-target soil micro-organisms and their ecosystem function when used as directed.


Innovation. Quality. Solutions.

Zampro® Fungicide

Zampro Summary

FEATURE	ADVANTAGE	BENEFIT
New mode of action (Group 45)	Qxl site of action = no cross-resistance	Disease resistance management
Locally systemic activity (Group 40)	Translaminar movement	Protects upper and lower leaf surfaces
Active on downy mildew (Groups 40+45)	Reduces pressure on following products	Excellent for fungicide rotation
Quick adsorptive / absorption onto plants	Excellent rainfastness	Spray is not washed off if rain occurs
Widely compatible	Mix with powdery mildew sprays	Increased application efficiency.

VINEYARD PRODUCTS TIMELINE



- Note:**
- WHP: Dimethomorph application to grapes grown for export wine production in Australia**
Zampro or other products containing dimethomorph can be applied up to growth stage E-L 31 (berries pea-size 7 mm in diameter) provided that only ONE application is made per crop of ANY of these products. If more than one application is to be made, the current E-L 25 (80% capfall) restriction applies in order to avoid a detectable residue in wine.
 - Do NOT apply Surround to table grapes after fruit set.**
 - Do NOT use Fetrilon Combi 2 or Wuxal after E-L 20 on table grapes.**

The information provided herein may include extracts from the product label and does not constitute the complete directions for use. READ THE PRODUCT LABEL THOROUGHLY BEFORE OPENING OR USING ZAMPRO

AgNova Technologies Pty Ltd
ABN 70 097 705 158
PO Box 2069 Box Hill North
Victoria 3129 Australia
Ph 03 9899 8100

Data referred to herein were generated using products and rates that were registered at the time. Always read and follow product labels. AgNova Technologies Pty Ltd shall not be liable for any consequential or other loss or damage relating to the supply or subsequent handling or use of this product, unless such liability by law cannot be lawfully excluded or limited. All warranties, conditions or rights implied by statute or other law which may be lawfully excluded are so excluded. Where the liability of AgNova Technologies Pty Ltd for breach of any such statutory warranties and conditions cannot be lawfully excluded but may be limited to it re-supplying the product or an equivalent product or the cost of a product or an equivalent product, then the liability of AgNova Technologies Pty Ltd for any breach of such statutory warranty or condition is so limited.

© Copyright AgNova Technologies 2017. ZAM190913
® Zampro is a registered trademark of BASF. Kusabi is a registered trademark of Ishihara Sangyo Kaisha Ltd. Japan. Flute is a registered trademark of Nippon Soda Co. Ltd. Surround is a registered trademark of Tessenderlo Kerley, Inc. Wuxal is a registered trademark of AGLUKON Spezialdünger GmbH & Co. KG. Fetrilon is a registered trademark of COMPO GmbH & Co. KG. Metarex is a registered trademark of De Sangosse SA, France.
* Registered trademark.