

PEEL
HERE

CAUTION

KEEP OUT OF REACH OF CHILDREN

READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Flute[®] 50 EW
FUNGICIDE

ACTIVE CONSTITUENT: 50 g/L CYFLUFENAMID

GROUP U6 FUNGICIDE

A fungicide for the control of powdery mildew in grapevines and cucurbits as per the Directions for Use Table.

IMPORTANT: Read the attached booklet before using this product.

In a transport emergency dial 000, Police or Fire Brigade.

For specialist advice in an emergency only, call 1800 033 111 (24 hours).



AgNova Technologies Pty Ltd

ABN 70 097 705 158

Suite 3, 935 Station Street

Box Hill North, Vic. 3129

Australia

Phone (03) 9899 8100

agnova.com.au



DIRECTIONS FOR USE

RESTRAINTS

DO NOT apply by air as uneven coverage is likely to result.

DO NOT apply when wind speed is less than 3 or more than 20 km/h at the application site.

Grapevines

Except when applying with vineyard airblast equipment, **DO NOT** apply with spray droplets smaller than a **MEDIUM** spray droplet size category according to nozzle manufacturer specifications that refer to the ASAE S572 Standard or the British Crop Production Council guideline.

Cucurbits

DO NOT apply with spray droplets smaller than a **MEDIUM** spray droplet size category according to nozzle manufacturer specifications that refer to the ASAE S572 Standard or the British Crop Production Council guideline.

DO NOT apply during surface temperature inversion conditions at the application site.

Users of this product MUST make an accurate written record of the details of each spray application within 24 hours following application and KEEP this record for a minimum of 2 years. The spray application details that must be recorded are:

- 1. date with start and finish times of application;**
- 2. location address and paddock/s sprayed;**
- 3. full name of this product;**
- 4. amount of product used per hectare and number of hectares applied to;**
- 5. crop/situation and weed/pest;**
- 6. wind speed and direction during application;**
- 7. air temperature and humidity;**
- 8. nozzle brand, type, spray angle, nozzle capacity and spray system pressure measured during application;**
- 9. name and address of person applying this product. (Additional record details may be required by the state or territory where this product is used.)**

CROPS	DISEASE	RATE	WHP	CRITICAL COMMENTS
Grapevines	Powdery mildew (<i>Erysiphe necator</i>)	<i>Dilute spraying:</i> 35 mL/100 L water <i>Concentrate spraying:</i> Refer to the Mixing/ Application section	5 weeks	Apply before first sign of disease or when conditions (varietal and climatic) are conducive to disease development as part of a protectant disease management program. Apply sprays at 10 to 21 day intervals. Use the shorter intervals under conditions favouring disease infection, and at the start of the season (first leaf separated from shoot tip EL7) when there has been a heavy carry-over of powdery mildew infection, or during periods of rapid vine growth. Do not apply more than 2 (two) Flute® 50 EW sprays per season, and follow the CropLife Australia resistance management guidelines. Apply in sufficient water volume to obtain thorough coverage of the crop. Apply by dilute or concentrate spraying equipment. Apply the same total amount of product to the target crop whether applying by dilute or concentrate spraying method. Do not apply at concentrations greater than 3 X (i.e. concentrated volumes that require rates greater than 105 mL/100 L). The required dilute spray volume will change and the sprayer setup and operation may also need to be changed, as the crop grows.

CROPS	DISEASE	RATE	WHP	CRITICAL COMMENTS
Cucurbits	Powdery mildew (<i>Podosphaera xanthii</i>)	250 mL/ha or <i>Glasshouse:</i> 25 mL/100 L	1 day	Apply before first sign of disease or when conditions are conducive to disease development as part of a disease management program. Apply sprays at 7 to 10 day intervals. Use the shorter intervals under conditions favouring disease infection. <i>Glasshouse:</i> To ensure thorough spray coverage of the target crop, apply to the point of run-off. The required dilute spray volume will change and the sprayer setup and operation may also need to be changed, as the crop grows. For resistance management: Do not apply FLUTE 50 EW as consecutive sprays or apply more than 2 sprays per crop.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIOD

GRAPES (DOMESTIC): DO NOT HARVEST FOR 5 WEEKS AFTER APPLICATION

GRAPES (FOR WINE INTENDED FOR EXPORT): CONTACT AGNOVA TECHNOLOGIES PTY LTD, OR CONSULT YOUR WINERY OR THE AUSTRALIAN WINE RESEARCH INSTITUTE (AWRI) FOR THE RECOMMENDED WITHHOLDING PERIOD

CUCURBITS: DO NOT HARVEST FOR 1 DAY AFTER APPLICATION

EXPORT OF TREATED PRODUCE

Growers should note that MRLs or import tolerances do not exist in all markets for fruit treated with FLUTE 50 EW.

Fruit: If growing fruit for export (either fresh or dried), consult your peak industry body or AgNova Technologies Pty Ltd for the latest information on MRLs and import tolerances.

Wine: FLUTE 50 EW should not be applied later than EL31 [berries pea-size] when grapes are to be used to make wine for export. Contact your winery or the Australian Wine Research Institute (AWRI) at www.awri.com.au for the updated information on MRLs and overseas import tolerances BEFORE using FLUTE 50 EW.

GENERAL INSTRUCTIONS

MIXING/APPLICATION

Add the required quantity of FLUTE 50 EW to a partly filled spray tank and agitate. Complete filling while agitating. Ensure the spray mixture is properly agitated before restarting after stoppage.

Grapevines

Dilute Spraying

- Use a sprayer designed to apply high volumes of water up to the point of run-off and matched to the crop being sprayed.
- Set up and operate the sprayer to achieve even coverage throughout the crop canopy. Apply sufficient water to cover the crop to the point of run-off. Avoid excessive run-off.
- The required water volume may be determined by applying different test volumes, using different settings on the sprayer, from industry guidelines or expert advice.
- Add the amount of product specified in the Directions for Use table for each 100 L of water. Spray to the point of run-off.
- The required dilute spray volume will change and the sprayer setup and operation may also need to be changed as the crop grows.

Concentrate Spraying

- Use a sprayer designed and set up for concentrate spraying (that is a sprayer which applies water volumes less than those required to reach the point of run-off) and matched to the crop being sprayed.
- Set up and operate the sprayer to achieve even coverage throughout the crop canopy using your chosen water volume.
- Determine an appropriate dilute spray volume (see *Dilute Spraying* above) for the crop canopy. This is needed to calculate the concentrate mixing rate.
- The mixing rate for concentrate spraying can then be calculated in the following way:

EXAMPLE ONLY

1. Dilute spray volume as determined above: For example 1500 L/ha
 2. Your chosen concentrate spray volume: For example 500 L/ha
 3. The concentration factor in this example is: 3 X (i.e. $1500 \text{ L} \div 500 \text{ L} = 3$)
 4. If the dilute label rate is 35 mL/100 L, then the concentrate rate becomes 3 x 35, that is 105 mL/100 L of concentrate spray.
- The chosen spray volume, amount of product /100 L of water, and the sprayer setup and operation may need to be changed as the crop grows.
 - Do not use a concentration rate higher than that specified in the Critical Comments.
 - Use a spray volume of at least 200 L/ha.
 - For further information on concentrate spraying, users are advised to consult relevant industry guidelines, undertake appropriate competency training and follow industry Best Practices.

FUNGICIDE RESISTANCE WARNING **GROUP U6 FUNGICIDE**

FLUTE 50 EW is a member of the amidoxime group of fungicides. For fungicide resistance management, the product is a Group U6 fungicide. Some naturally occurring individual fungi resistant to FLUTE 50 EW and other Group U6 fungicides may exist through normal genetic variability in any fungal population. The resistant individuals can eventually dominate the fungal population if these fungicides are used repeatedly. These resistant fungi will not be controlled by FLUTE 50 EW or other Group U6 fungicides, thus resulting in a reduction in efficacy and possible yield loss.

Since the occurrence of resistant fungi is difficult to detect prior to use, AgNova Technologies Pty Ltd accepts no liability for any losses that may result from the failure of FLUTE 50 EW to control resistant fungi. Refer to Directions for Use table and published CropLife Resistance Management Strategies www.croplifeaustralia.org.au.

COMPATIBILITY

For information on compatibility contact your local AgNova representative or visit our website at agnova.com.au

RE-ENTRY PERIOD

Do not allow entry into treated areas until the spray has dried, unless wearing cotton overalls buttoned to the neck and wrist (or equivalent clothing) and elbow-length chemical resistant gloves. Clothing must be laundered after each day's use. For glasshouses and other enclosed areas, do not allow re-entry until spray has dried or dissipated, and enclosed area has been thoroughly ventilated.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

DO NOT apply under weather conditions or from spraying equipment that may cause spray to drift onto nearby susceptible plants/crops, cropping lands or pastures.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

Very toxic to aquatic life. DO NOT contaminate wetlands or water courses with this product or used containers.

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight.

Triple rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point.

If not recycling, break, crush or puncture and deliver empty packaging to an approved waste management facility. DO NOT burn empty containers or product.

If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations.

SAFETY DIRECTIONS

Will irritate the skin. May irritate the eyes. Avoid contact with eyes and skin. When opening the container, preparing spray and using the prepared spray, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing) and elbow-length chemical resistant gloves. If product on skin, immediately wash area with soap and water. Wash hands after use. After each day's use, wash gloves and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26

SAFETY DATA SHEET

If additional hazard information is required refer to the Material Safety Data Sheet. For a copy visit our website at agnova.com.au

CONDITIONS OF SALE

AgNova Technologies Pty Ltd shall not be liable for any consequential or other loss or damage relating to the supply or subsequent handling or use of this product, unless such liability by law cannot be lawfully excluded or limited. All warranties, conditions or rights implied by statute or other law which may be lawfully excluded are so excluded. Where the liability of AgNova Technologies Pty Ltd for breach of any such statutory warranties and conditions cannot be lawfully excluded but may be limited to it re-supplying the product or an equivalent product or the cost of a product or an equivalent product, then the liability of AgNova Technologies Pty Ltd for any breach of such statutory warranty or condition is so limited.

In a transport emergency dial 000, Police or Fire Brigade.
For specialist advice in an emergency only, call
1800 033 111 (24 hours).

AgNova Technologies Pty Ltd

ABN 70 097 705 158

Suite 3, 935 Station Street

Box Hill North, Vic. 3129 Australia

Phone (03) 9899 8100

agnova.com.au

Batch Number

Date of Manufacture

APVMA Approval No: 66970/100670

® Registered Trademark of Nippon Soda Co., Ltd.

