



FRUITITION[®]

FRUIT FLY LURE & TRAP
FOR QUEENSLAND FRUIT FLY (*Bactrocera tryoni*)

USE INSTRUCTIONS & **ASSEMBLY GUIDE**

NZ Patent No. 630903 and patents pending.

FRUITION®

The Revolutionary Lure & Trap for Qld Fruit Fly

Fruition Traps are a unique new system for monitoring and managing fruit fly populations, and were developed specifically to target egg-laying females.

Fruition Traps have proven to be highly effective in attracting Queensland fruit fly (*Bactrocera tryoni*) through a unique combination of colour, shape and smell.

Fruit flies are drawn to the vicinity of the trap using aromas that resemble those of ripe fruit, and once in visual range, are able to detect the colour and shape of the trap structure. A sticky surface then traps the flies in such a way that they can be easily identified and counted.

As soon as fruit flies are detected, growers should implement a full Integrated Pest Management (IPM) control program.

HOW TO USE FRUITION TRAPS

Fruition Traps can be used for both population monitoring and control as part of an IPM control program, when susceptible crops are fruiting.

FOR FRUIT FLY MONITORING

Fruition Traps are suitable for all crops where there is a need to monitor for the presence of egg-laying fruit flies before crop damage occurs.

Fruition Traps should be installed at a rate of 15 per hectare from the early stages of fruit set. Traps should be monitored regularly, preferably on a daily basis, with records maintained for each monitoring event.

As soon as fruit flies are detected, growers should implement a full IPM control program (see below). Regular monitoring of traps should continue throughout the season to ensure that the control program is adequate.

FOR FRUIT FLY CONTROL

Efficacy of a fruit fly control program is dependent on a range of factors including pest pressure during the season. For effective management of fruit fly, Fruition Traps should be used as part of a broader strategic control program, involving other products approved for the control of fruit fly, as described below.

As soon as fruit flies are detected on the monitoring traps, commence an IPM control program as follows:

1. Increase the number of Fruition Traps based on crop susceptibility. As a minimum, use;
 - a. Low susceptibility crops: 15-30 traps per hectare
 - b. Moderate to High susceptibility crops: 30-50 traps per hectare
2. Trap numbers may need to be increased above the minimum based on a range of factors including fruit fly numbers trapped during the monitoring phase, crop history & susceptibility, crop canopy size & density, crop value and seasonal conditions, etc.
3. Commence applications of a premium quality gelatinised yeast bait according to the label and reapply at least every seven (7) days. A registered insecticide should be included with the bait according to the insecticide label.
4. Continue monitoring for the presence of fruit flies until immediately after final harvest.
5. If fruit fly numbers on Fruition Traps indicate high or erratic pest pressure as fruit develops and becomes more susceptible to fruit fly, insecticide control options should be included in the IPM control program.
6. Traps should be replaced if;
 - a. sticky surfaces are heavily covered by fruit flies or foreign objects,
 - b. the lure sachet has expired,
 - c. damaged or missing.

Typically, an effective IPM control program will result in the number of freshly trapped fruit flies declining over time.



PLACEMENT OF FRUITION TRAPS

Fruition Traps should be placed evenly around and throughout the site.

Traps should be hung in the fruit zone, usually 1.5 to 2 metres above the ground. Ideally traps will be in the tree canopy in a location away from surrounding branches and clearly visible within the orchard.

In vegetable crops, traps should be hung immediately above the crop and in adjacent trees/vegetation within 5 metres of the crop.

Ideally neighbouring crops will also be monitored as these can be a source of fruit fly populations.

HOW LONG DO FRUITION TRAPS LAST?

The Fruition Trap gel lure in the open sachet will continue to operate for a period of up to eight (8) weeks, during which time any remaining gel will change colour from blue to very pale blue or white, signifying the need to replace the entire trap.

Traps may need replacing sooner if;

- a. sticky surfaces are heavily covered by fruit flies or foreign objects,
- b. the lure sachet has expired,
- c. damaged or missing.



STORAGE AND DISPOSAL OF FRUITION TRAPS

Store unused Fruition Traps and lure sachets in original packaging out of direct sunlight and below 25°C.

Keep out of reach of children and animals.

Dispose of the used traps at the end of the cropping season through a waste disposal system in compliance with local, state or territory government regulations.

HANDLING PRECAUTIONS

Avoid contact with the adhesive surface. The adhesive is non-toxic and vegetable oil can be used to remove from skin.

SAFETY DATA SHEET

If additional information is required refer to the Safety Data Sheet. For a copy visit our website agnova.com.au









CONDITIONS OF SALE

AgNova Technologies Pty Ltd shall not be liable for any consequential or other loss or damage relating to the supply or subsequent handling or use of this product, unless such liability by law cannot be lawfully excluded or limited. All warranties, conditions or rights implied by statute or other law which may be lawfully excluded are so excluded. Where the liability of AgNova Technologies Pty Ltd for breach of any such statutory warranties and conditions cannot be lawfully excluded but may be limited to it re-supplying the product or an equivalent product or the cost of a product or an equivalent product, then the liability of AgNova Technologies Pty Ltd for any breach of such statutory warranty or condition is so limited.

This product does not require APVMA approval.

Fruition Trap Assembly Guide

Each trap consists of two discs, a lure sachet, a metal clip and a cable tie. They are supplied with the discs loosely fitted together and the clip tightly fitted to one disc.



AgNova Technologies Pty Ltd
ABN 70 097 705 158
PO Box 2069 Box Hill North
Victoria 3129 Australia
Ph 03 9899 8100

AgNova Technologies Pty Ltd shall not be liable for any consequential or other loss or damage relating to the supply or subsequent handling or use of this product, unless such liability by law cannot be lawfully excluded or limited. All warranties, conditions or rights implied by statute or other law which may be lawfully excluded are so excluded. Where the liability of AgNova Technologies Pty Ltd for breach of any such statutory warranties and conditions cannot be lawfully excluded but may be limited to it re-supplying the product or an equivalent product or the cost of a product or an equivalent product, then the liability of AgNova Technologies Pty Ltd for any breach of such statutory warranty or condition is so limited.

* Fruition is a registered trademark of AgNova Technologies Pty Ltd

© Copyright AgNova Technologies 2016. FRULAB161102