

READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Lorox[®]
Linuron DF
herbicide

ACTIVE CONSTITUENT: 500 g/kg LINURON

GROUP 5 HERBICIDE

For selective weed control in carrots, parsnips, onions, leeks, shallots & spring onions, potato, celery, sweet corn & maize, pyrethrum, celeriac, soybeans, cereals and coriander seed crops as per the Directions for Use table.

Distributed by:

AgNova Technologies Pty Ltd
ABN 70 097 705 158
Unit 4, 482 Kingsford Smith Drive
Hamilton, Qld 4007 Australia
Phone (03) 9899 8100
Email info@agnova.com.au
agnova.com.au



Manufactured for:
Tessengerlo Kerley, Inc.
2910 N. 44th Street, Suite 100
Phoenix, Arizona 85018 USA



An American Vanguard Company

Net Contents 2 kg and 10 kg

DIRECTIONS FOR USE

For use in all States unless indicated otherwise.

RESTRAINTS

Sufficient moisture, either by irrigation or rainfall, is required within 3–4 days of application to carry the product into the soil profile.

DO NOT add wetting agents, surfactants or crop oils unless specified in Critical Comments.

DO NOT apply to weeds under severe stress due to, for example, very dry or windy conditions, disease, etc.

DO NOT allow mixture to stand in vat for prolonged periods, as it is difficult to re-suspend and may lose effectiveness.

DO NOT apply with misting machines.

SPRAY DRIFT RESTRAINTS

Specific definitions for terms used in this section of the label can be found at apvma.gov.au/spraydrift

DO NOT allow bystanders to come into contact with the spray cloud.

DO NOT apply in a manner that may cause an unacceptable impact to native vegetation, agricultural crops, landscaped gardens and aquaculture production, or cause contamination of plant or livestock commodities, outside the application site from spray drift.

Wherever possible, correctly use application equipment designed to reduce spray drift and apply when the wind direction is away from these sensitive areas.

DO NOT apply unless the wind speed is between 3 and 20 kilometres per hour at the application site during the time of application.

DO NOT apply if there are hazardous surface temperature inversion conditions present at the application site during the time of application. Surface temperature inversion conditions exist most evenings one to two hours before sunset and persist until one to two hours after sunrise.

CROP	WEEDS CONTROLLED	WEED STAGE	RATE/HA	CRITICAL COMMENTS
Carrots	Annual Broadleaved Weeds and Grasses such as: Amaranth (inc. Dwarf, Green, Slim, Red Root, Powell's), Blackberry Nightshade, Capeweed, Chickweed, Common Sowthistle, Corn Gromwell, Corn Spurry, Deadnettle, Doublegee/Three-cornered Jack (Spiny Emex), Fat-hen, Hedge Mustards, Heliotrope, Marsh Mallow, Mustard, Paterson's Curse, Pigweed, Potato Weed, Redshank, Rough Poppy, Scarlet Pimpernel, Shepherd's Purse, Slender Thistle, Staggerweed, Stinging Nettle, Summer Grass (Summer Grass not controlled in Qld or November to February in other states), Swinecress, Turnip Weed, Variegated Thistle, Wild Mustard, Wild Radish, Wild Turnip, Winter Grass	Pre-emergence	2.2 to 4.5 kg	Pre-emergence: Apply immediately after planting but before crop emerges. Use low rate on lighter soils and higher rates on heavier soils. DO NOT use on light sandy soils, low in clay or organic matter. Heavy rains after application may cause crop damage. DO NOT replant treated areas to other crops within 3 months of application.
		Post-emergence Weeds no more than 7 cm tall	1.1 to 2.2 kg	Post-emergence: DO NOT apply before carrots have reached the 4 true leaf stage. DO NOT apply within 5 days of any application of any emulsifiable pesticide or wetting agent as damage to crop may result. Use low rate on lighter soils or early crop development stage and higher rates on heavier soils or later crop development stages. On soils high in organic matter use post-emergence rather than pre-emergence. Heavy rains after application may cause crop damage. DO NOT replant treated areas to other crops within 3 months of application.

CROP	WEEDS CONTROLLED	WEED STAGE	RATE/HA	CRITICAL COMMENTS
Parsnips	Annual Broadleaved Weeds and Grasses such as: Amaranth (inc. Dwarf, Green, Slim, Red Root, Powell's), Blackberry Nightshade, Capeweed, Chickweed, Common Sowthistle, Corn Gromwell, Corn Spurry, Deadnettle, Doublegee/Three-cornered Jack (Spiny Emex), Fat-hen, Hedge Mustards, Heliotrope, Marsh Mallow, Mustard, Paterson's Curse, Pigweed, Potato Weed, Redshank, Rough Poppy, Scarlet Pimpernel, Shepherd's Purse, Slender Thistle, Staggerweed, Stinging Nettle, Summer Grass (Summer Grass not controlled in Qld or November to February in other states), Swinecress, Turnip Weed, Variegated Thistle, Wild Mustard, Wild Radish, Wild Turnip, Winter Grass	Pre-emergence	2.2 to 4.5 kg	Pre-emergence: Apply immediately after planting but before crop emerges. Use low rates on lighter soils and higher rates on heavier soils. DO NOT use on light sandy soils, low in clay or organic matter. Heavy rains after application may cause crop damage. DO NOT replant treated areas to other crops within 3 months of application.
	Annual Broadleaved Weeds such as: Amaranth (inc. Dwarf, Green, Slim, Red Root, Powell's), Blackberry Nightshade, Capeweed, Chickweed, Common Sowthistle, Corn Gromwell, Corn Spurry, Deadnettle, Doublegee/Three-cornered Jack (Spiny Emex), Fat-hen, Hedge Mustards, Heliotrope, Marsh Mallow, Mustard, Paterson's Curse, Pigweed, Potato Weed, Redshank, Rough Poppy, Scarlet Pimpernel, Shepherd's Purse, Slender Thistle, Staggerweed, Stinging Nettle, Swinecress, Turnip Weed, Variegated Thistle, Wild Mustard, Wild Radish, Wild Turnip	Post-emergence Early weed post-emergence from the 1–2 true leaf stage	200 to 700 g	Post-emergence: DO NOT apply before parsnips are in 2–3 true leaf stage. Use the lower rates on light soils or early crop development stage, and the higher rates on heavier soils or later crop development stages. Apply as a foliar application by boomspray in spray volume of 500 L/ha. DO NOT apply more than one application per crop. DO NOT apply if temperature is above 30°C. DO NOT apply to diseased or stressed crops. The sensitivity of parsnips has not been fully evaluated. Treat a small area to ascertain the reaction before treating the whole crop. Heavy rains after application may cause crop damage.
Onions	Annual Broadleaved Weeds such as: Amaranth (inc. Dwarf, Green, Slim, Red Root, Powell's), Black Bindweed, Capeweed, Charlock, Chickweed, Clover, Common Sowthistle, Corn Gromwell, Corn Spurry, Deadnettle, Doublegee/Three-cornered Jack (Spiny Emex), Fat-hen, Field Cress, Hedge Mustards, Lesser Swinecress, Medics, Mustard, Pigweed, Prickly Sowthistle, Rough Poppy, Shepherd's Purse, Slender Thistle, Spear Thistle, Stinging Nettle, Turnip Weed, Wild Mustard, Wild Radish, Wild Turnip	Post-emergence	300 to 550 g (Qld, NSW, Vic, SA, WA only)	Apply when onions are 15 cm or more in height. Retreat when necessary. Use lower rates on light soils or early crop development stages, and the higher rate on heavier soils.
			700 to 1 kg (Tas only)	Apply lowest rate when onion plants are 2–3 leaf stage and high rate from 3 leaf stage onwards. Up to three applications may be used.

CROP	WEEDS CONTROLLED	WEED STAGE	RATE/HA	CRITICAL COMMENTS
Leeks	Annual Broadleaved Weeds such as: Amaranth (inc. Dwarf, Green, Slim, Red Root, Powell's), Black Bindweed, Capeweed, Charlock, Chickweed, Clover, Common Sowthistle, Corn Gromwell, Corn Spurry, Deadnettle, Doublegee/Three-cornered Jack (Spiny Emex), Fat-hen, Field Cress, Hedge Mustards, Lesser Swinecress, Medics, Mustard, Pigweed, Prickly Sowthistle, Rough Poppy, Shepherd's Purse, Slender Thistle, Spear Thistle, Stinging Nettle, Turnip Weed, Wild Mustard, Wild Radish, Wild Turnip	Post-emergence	300 to 500 g	<p>Apply when leeks are 15 cm or more in height. Apply as required (5–10 weeks after transplanting is common), over the top of the crop, using a boom spray. Can be applied over seeded crops at the same stage.</p> <p>DO NOT apply more than 3 applications per crop.</p> <p>If weed control is less than adequate and resistance is suspected, the second application should not be applied.</p> <p>Post application, leeks may display slight tip burn, and slow the growth of the crop. These issues are transient. Ensure the crop has recovered before sequential applications. Limit first use to small areas before wider application to the whole crop to ensure Lorox® Linuron DF is suitable to the variety as well as the site's soil & environmental conditions.</p> <p>DO NOT apply within 5 days of any application of an emulsifiable pesticide or wetting agent as damage to crop may result.</p> <p>Injury caused by sand blasting, water stress or other causes may render leeks sensitive to Lorox Linuron DF and lead to unacceptable crop damage.</p>
Shallots and Spring Onions	Annual Broadleaved Weeds such as: Amaranth (inc. Dwarf, Green, Slim, Red Root, Powell's), Black Bindweed, Capeweed, Charlock, Chickweed, Clover, Common Sowthistle, Corn Gromwell, Corn Spurry, Deadnettle, Doublegee/Three-cornered Jack (Spiny Emex), Fat-hen, Field Cress, Hedge Mustards, Lesser Swinecress, Medics, Mustard, Pigweed, Prickly Sowthistle, Rough Poppy, Shepherd's Purse, Slender Thistle, Spear Thistle, Stinging Nettle, Turnip Weed, Wild Mustard, Wild Radish, Wild Turnip	Post-emergence	300 to 500 g	<p>Apply after 3 true leaf stage, when shallots and spring onions are 15 cm high. Foliage must be waxy.</p> <p>Apply to foliage by aerial or ground boomspray or equivalent technology.</p> <p>DO NOT apply more than 2 applications per crop.</p> <p>DO NOT apply less than 14–21 days after the initial treatment.</p> <p>Use lower rates on light soil or on earlier crop growth stages, and the higher rates can be applied on heavier soils. Application can cause some tip burn or leaf bending that may slow the crop's growth, however this is likely to be transient.</p> <p>Injury caused by sand blasting, water stress or other causes may render spring onions or shallots sensitive to Lorox Linuron DF and lead to unacceptable crop damage.</p> <p>Limit first use to small areas before wider application to the whole crop to ensure Lorox Linuron DF is suitable to the variety as well as the site's soil & environmental conditions.</p> <p>DO NOT apply within 5 days of any application of an emulsifiable pesticide or wetting agent as damage to crop may result.</p>
Potatoes	Annual Broadleaved Weeds and Grasses such as: Amaranth (inc. Dwarf, Green, Slim, Red Root, Powell's), Apple-of-Peru, Blackberry Nightshade, Capeweed, Chickweed, Clover, Common Sowthistle, Corn Gromwell, Corn Spurry, Deadnettle, Doublegee/Three-cornered Jack (Spiny Emex), Fat-hen, Hedge Mustards, Mustard, Pigweed, Potato Weed, Redshank, Rough Poppy, Shepherd's Purse, Slender Thistle, Staggerweed, Stinging Nettle, Turnip Weed, Variegated Thistle, Wild Mustard, Wild Radish, Wild Turnip, Winter Grass, Winter Weed, Wireweed	Pre-emergence	2.2 to 4.5 kg	<p>Pre-emergence: Apply after planting but before the emergence of crop. Incorporate by irrigation or rainfall as soon as possible after application. DO NOT cultivate after application. Use higher rate during summer months.</p> <p>Use low rates on lighter soils and higher rates on heavier soils.</p>

CROP	WEEDS CONTROLLED	WEED STAGE	RATE/HA		CRITICAL COMMENTS
Celery	Annual Broadleaved Weeds and Grasses such as: Amaranth (inc. Dwarf, Green, Slim, Red Root, Powell's), Blackberry Nightshade, Capeweed, Chickweed, Common Sowthistle, Corn Gromwell, Corn Spurry, Deadnettle, Doublegee/Three-cornered Jack (Spiny Emex), Fat-hen, Hedge Mustards, Heliotrope, Marsh Mallow, Mustard, Paterson's Curse, Pigweed, Potato Weed, Redshank, Rough Poppy, Scarlet Pimpernel, Shepherd's Purse, Slender Thistle, Staggerweed, Stinging Nettle, Summer Grass (Summer Grass not controlled in Qld or November to February in other states), Swinecress, Turnip Weed, Variegated Thistle, Wild Mustard, Wild Radish, Wild Turnip, Winter Grass	Post-emergence Apply at the 2-4 true leaf stage Weeds no more than 7 cm tall	1 kg		Apply once transplants are established, as soon as new growth has started. DO NOT apply after the 5 leaf stage of the crop. DO NOT apply if temperature exceeds 30°C. DO NOT apply within 5 days of any application of an emulsifiable pesticide or wetting agent as damage to crop may result. The sensitivity of celery has not been fully evaluated. Treat a small area to ascertain the reaction before treating the whole crop.
Sweet Corn & Maize	Apply to Annual Broadleaved Weeds and Grasses such as: Amaranth (inc. Dwarf, Green, Slim, Red Root, Powell's), Apple-of-Peru, Barnyard Grass, Bathurst Burr, Black Bindweed, Blackberry Nightshade, Capeweed, Common Sowthistle, Corn Gromwell, Deadnettle, Doublegee/Three-cornered Jack (Spiny Emex), Fat-hen, Hedge Mustards, Mustard, Noogoora Burr, Pigweed, Potato Weed, Redshank, Stinging Nettle, Stinking Roger, Summer Grass, Turnip Weed, Wild Mustard, Wild Radish, Wild Turnip Apply where grasses are not a problem.	Pre-emergence	-	2.2 to 3.5 kg on heavy soils only	Pre-emergence: Apply as a band of broadcast treatment immediately after planting. DO NOT use on light sandy soils low in clay or organic matter. Heavy rains after application may cause crop damage.
	Refer to weeds above. Apply where Barnyard Grass or Summer Grass are a problem.		-	4.5 to 5.5 kg on heavy soils only	
	Refer to weeds above. Apply where grasses are not a problem.	Post-emergence	1.1 to 2.2 kg on light soils	2.2 to 4.5 kg on heavy soils	Post-emergence: Apply as directed band spray to each side row when crop is 30-40 cm high. Avoid spray contact with crop foliage as injury may result. DO NOT replant treated areas to other crops within 3 months of application. If weeds are present add a wetting agent at the use rate on its label.
	Refer to weeds above. Apply where Barnyard Grass or Summer Grass are a problem.		1.1 to 3.5 kg on light soils	2.2 to 5.5 kg on heavy soils	
Pyrethrum	Lesser Swinecress, Wild Radish, Shepherd's Purse, Volunteer Potato, Chickweed, Wild Turnip	Post-emergence only Apply when weeds are still in the seedling stage	180 to 450 g		Apply from 4 true leaf stage of seedling pyrethrum. Applications to be made in the crop's establishment year, prior to the first cut. Avoid application to stressed plants or applications during the hottest part of the day. The sensitivity of pyrethrum has not been fully evaluated. Treat a small area of crop and monitor plant health closely over the following days/ weeks. Discontinue applications when any adverse symptoms post-treatment are observed.
	Clover		300 to 450 g		

CROP	WEEDS CONTROLLED	WEED STAGE	RATE/HA	CRITICAL COMMENTS
Celeriac (trans-planted)	Annual Broadleaved Weeds and Grasses such as: Amaranth (inc. Dwarf, Green, Slim, Red Root, Powell's), Blackberry Nightshade, Capeweed, Chickweed, Common Sowthistle, Corn Gromwell, Corn Spurry, Deadnettle, Doublegee/Three-cornered Jack (Spiny Emex), Fat-hen, Hedge Mustards, Heliotrope, Marsh Mallow, Mustard, Paterson's Curse, Pigweed, Potato Weed, Redshank, Rough Poppy, Scarlet Pimpernel, Shepherd's Purse, Slender Thistle, Staggerweed, Stinging Nettle, Summer Grass (Summer Grass not controlled in Qld or November to February in other states), Swinecress, Turnip Weed, Variegated Thistle, Wild Mustard, Wild Radish, Wild Turnip, Winter Grass	Post-emergence Apply at the 2–4 true leaf stage Ensure weeds are no more than 7 cm tall	1.1 to 2.2 kg	Apply once transplants are established, as soon as new growth has started. DO NOT apply after the 5 leaf stage of the crop. DO NOT apply more than one (1) application per crop. DO NOT apply within 5 days of any application of an emulsifiable pesticide or wetting agent as damage to crop may result. Use the lower rates on light sandy soils, and higher rates on heavier soils or when longer residual control is desired. DO NOT apply if temperature exceeds 30°C.
Soybeans	Annual Broadleaved Weeds and Grasses such as: Amaranth (inc. Dwarf, Green, Slim, Red Root, Powell's), Apple-of-Peru, Barnyard Grass, Bathurst Burr, Black Bindweed, Blackberry Nightshade, Capeweed, Common Sowthistle, Corn Gromwell, Doublegee/ Three-cornered Jack (Spiny Emex), Fat-hen, Hedge Mustards, Mustard, Noogoora Burr, Pigweed, Potato Weed, Redshank, Rough Poppy, Shepherd's Purse, Slender Thistle, Stinking Roger, Summer Grass, Turnip Weed, Wild Radish, Wild Turnip	Pre-emergence	2.2 to 5.5 kg (Qld, NSW, Vic, WA only)	Pre-emergence: Apply within 2–3 days after sowing. Sow beans to a depth of at least 2.5 cm. Heavy rains after application may cause crop damage. Use lower rates on lighter soils, and higher rates on heavier soils. DO NOT replant treated area to other crops within 3 months of application.
Wheat, Oats & Barley	Capeweed, Charlock, Corn Gromwell (Sheepweed, White Ironweed), Deadnettle, Doublegee/Three-cornered Jack (Spiny Emex), Fat-hen, Hedge Mustards, Indian Hedge Mustard, Mustard, Rough Poppy, Shepherd's Purse, Slender Thistle, Stinging Nettle, Turnip Weed, Wild Radish, Wild Turnip	Post-emergence	500 g	For All Weeds: For best results apply early. DO NOT apply Lorox Linuron DF if weeds are greater than 15 cm in height or diameter, or crop is beyond early tillering or crop and weeds are under drought stress. Apply when weeds are less than 5 cm in height or diameter, crop is in 2–4 leaf stage and soil is moist enough for good growth of crop and weed seedlings.
	As above plus Amsinckia and Saffron Thistle		850 g	Apply if weeds are 5–15 cm in height or diameter and crop is in four leaf to early tillering stage.
Coriander Seed Crops	Amsinckia, Capeweed, Charlock, Corn Gromwell (Sheepweed, White Ironweed), Deadnettle, Doublegee/Three-cornered Jack (SpinyEmex), Fat-hen, Hedge Mustard, Indian Hedge Mustard, Mustard, Rough Poppy, Saffron Thistle, Slender Thistle, Stinging Nettle, Turnip Weed, Wild Radish, Wild Turnip	Pre-emergence	750 g to 1.0 kg (SA, Vic, NSW,Tas only)	DO NOT use this product on fresh coriander production. Use lower rate quoted on light soils and increase up to the higher rate for heavy clay soils or soils high in organic matter. Pre-emergence only: Apply immediately after sowing.
		Post-emergence		Post-emergence only: Apply from 2 leaf onwards. Higher rates give longer residual control. Wet soil and/or rain is a benefit soon after application.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIODS

HARVEST

PYRETHRUM – DO NOT HARVEST FOR 25 WEEKS AFTER APPLICATION.

CARROTS, CORIANDER, LEEKS, PARSNIPS (Post-emergence) – DO NOT HARVEST FOR 10 WEEKS AFTER APPLICATION.

CELERIAC – DO NOT HARVEST FOR 4 WEEKS AFTER APPLICATION.

BARLEY, CELERY, MAIZE, OATS, ONIONS, PARSNIPS (Pre-emergence), POTATOES, SHALLOTS, SPRING ONIONS, SOYBEANS, SWEET CORN, WHEAT – NOT REQUIRED WHEN USED AS DIRECTED.

GRAZING

PARSNIPS, SHALLOTS, SPRING ONIONS – DO NOT GRAZE OR CUT FOR STOCK FOOD.

TRADE ADVICE

Growers should note that maximum residue limits (MRLs) or import tolerances may not exist in all markets for edible produce treated with Lorox Linuron DF herbicide. If you are growing edible produce for export, please check with AgNova Technologies for the latest information on MRLs and import tolerances before using Lorox Linuron DF herbicide.

GENERAL INSTRUCTIONS

Lorox Linuron DF herbicide can be applied pre- and post-emergence to control certain annual grasses, broad-leaved weeds and some seedling perennial weeds in carrots, parsnips, onions, leeks, shallots & spring onions, potato, celery, sweet corn & maize, pyrethrum, celeriac, soybeans, cereals and coriander seed crops.

Lorox Linuron DF herbicide is a selective systemic herbicide, absorbed principally by the roots but also by the foliage, with translocation primarily acropetally in the xylem. Read the Directions for Use before using this product. Shake well before opening. For best results, apply to a well cultivated, weed-free surface which is free of trash and clods. DO NOT cultivate after application of the product as weed control may be reduced and/or crop injury may result.

Application

Lorox Linuron DF herbicide is a dry flowable formulation to be mixed with water and applied as a spray. Partially fill the spray tank with water. Measure the amount of product required as indicated in the directions for use table. Add this to the spray tank with the agitation system engaged. Top up tank to the correct water volume. To prevent nozzle blockage, strainer and nozzle screens should be 50 mesh or coarser.

THE MATERIAL MUST BE KEPT IN SUSPENSION AT ALL TIMES BY CONTINUOUS AGITATION AS IT IS DIFFICULT TO RE-SUSPEND AND MAY LOSE EFFECTIVENESS.

Water Volume

Ground Application:

- Pre-Emergence: 300–500 L/ha
- Post-Emergence: Use minimum of 300 L/ha
- Band Treatments: Use proportionately less.

Aerial Application:

Use a minimum of 20 L/ha spray under optimum conditions otherwise a minimum of 30 L/ha.

Spray Equipment Hygiene

Before and after spraying Lorox Linuron DF herbicide, spray equipment including tanks, nozzles, filters and spraylines should be cleaned thoroughly with a commercially available tank and equipment cleaner such as All Clear® DS.

RESISTANT WEEDS WARNING

GROUP 5 HERBICIDE

Lorox Linuron DF herbicide is a member of the Urea group of herbicides. Lorox Linuron DF herbicide has the inhibitors of photosynthesis at photosystem II mode of action. For weed resistance management Lorox Linuron DF herbicide is a Group 5 herbicide.

Some naturally-occurring weed biotypes resistant to Lorox Linuron DF herbicide and other inhibitors of photosynthesis at photosystem II herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds may not be controlled by Lorox Linuron DF herbicide or other inhibitors of photosynthesis at photosystem II herbicides.

Since the occurrence of resistant weeds is difficult to detect prior to use, Tessengerlo Kerley, Inc. accepts no responsibility for any losses that may result from the failure of Lorox Linuron DF herbicide to control resistant weeds.

Strategies to minimise the risk of herbicide resistance are available. Consult your farm chemical supplier, consultants, local Department of Agriculture or Primary industries, or local AgNova Technologies representative.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

Product is slightly toxic to fish. DO NOT contaminate streams, rivers, waterways with chemical or used containers.

STORAGE AND DISPOSAL

Keep out of reach of children.

Store in the closed, original container in a dry, cool, well-ventilated area out of direct sunlight. Store in a locked room or place away from children, animals, food, feedstuffs, seed and fertilisers. Shake and empty contents into spray tank. Do not dispose of undiluted chemicals on site. Break, crush or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.

SAFETY DIRECTIONS

May irritate the eyes and skin. Avoid contact with the eyes and skin. DO NOT inhale dust or spray mist. If product on skin, immediately wash the area with soap and water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water.

FIRST AID

First aid is not generally required. If in doubt, contact a Poisons Information Centre (e.g. phone Australia 13 11 26, New Zealand 0800 764 766) or a doctor.

SAFETY DATA SHEET

If additional hazard information is required refer to the Safety Data Sheet. For a copy visit our website at agnova.com.au

CONDITIONS OF SALE

AgNova Technologies Pty Ltd and Tessenderlo Kerley, Inc. shall not be liable for any consequential or other loss or damage relating to the supply or subsequent handling or use of this product, unless such liability by law cannot be lawfully excluded or limited. All warranties, conditions or rights implied by statute or other law which may be lawfully excluded are so excluded. Where the liability of AgNova Technologies Pty Ltd and Tessenderlo Kerley, Inc. for breach of any such statutory warranties and conditions cannot be lawfully excluded but may be limited to it re-supplying the product or an equivalent product or the cost of a product or an equivalent product, then the liability of AgNova Technologies Pty Ltd and Tessenderlo Kerley, Inc. for any breach of such statutory warranty or condition is so limited.

®Lorox is a registered trademark of Tessenderlo Kerley, Inc.

®All Clear is a registered trademark of AgNova Technologies Pty Ltd

Additional statements as required by Safe Work Australia
in accordance with the Globally Harmonised System of Classification and Labelling (GHS).

May damage the unborn child, suspected of damaging fertility. Suspected of causing cancer.

May cause damage to organs through prolonged or repeated exposure.

Harmful if inhaled. May be harmful if swallowed. Very toxic to aquatic life with long lasting effects.

Obtain, read and follow all safety instructions before use. Wear protective gloves and protective clothing. Use only outdoors or in a well-ventilated area. IF exposed or concerned, get medical advice. Get medical help if you feel unwell.

IF INHALED: Remove person to fresh air and keep comfortable for breathing. Get medical help. IF SWALLOWED: Get medical help. Store locked up. Avoid release to the environment not in accordance with this label. Collect spillage.

In a transport emergency dial 000, Police or Fire Brigade.
For specialist advice in an emergency only, call 1800 033 111 (24 hours).

APVMA Approval No: 60222/136517

Distributed by:

AgNova Technologies Pty Ltd

ABN 70 097 705 158

Unit 4, 482 Kingsford Smith Drive

Hamilton, Qld 4007 Australia

Phone (03) 9899 8100

Email info@agnova.com.au

agnova.com.au

Manufactured for:

Tessenderlo Kerley, Inc.

2910 N. 44th Street, Suite 100

Phoenix, Arizona 85018 USA



An American Vanguard Company